

Olmesartan attenuates type 2 diabetes-associated liver injury: Cross-talk of AGE/RAGE/JNK, STAT3/SCOS3 and RAS signaling pathways

Hanan Salah ,Nesma M E Abo El-Nasr, Dalia Osama Saleh, Sawsan S Mahmoud, Salwa M Nofal, Rania M Abdelsalam, Marwa M Safar

Abstract

Olmesartan (OLM), an angiotensin receptor blocker, was tested against diabetes/insulin resistance (IR) models associated with renal/cardiovascular complications. Methods: we tested its potential role against diabetes-induced hepatic injuries using an IR/type2 diabetic (IR/D) model induced by high fat/high fructose diet for 7 weeks + a single sub-diabetogenic dose of streptozotocin (35mg/kg; i.p). IR/D rats were orally treated with OLM (10 mg/kg), pioglitazone (PIO; 5 or 10 mg/kg) or their combinations for 4 consecutive weeks. OLM alone opposed the detrimental effects of IR/D; it significantly improved metabolic parameters, liver function, and abated hepatic oxidative stress, and inflammatory cytokine interleukin-6 (IL-6) and its upstream mediator nuclear factor kappa B. Consequently, OLM turned off the downstream cue p-Jak2/STAT3/SOCS3. Moreover, it suppressed the elevated AGE/RAGE/p-JNK pathway and increased the PPAR γ adiponectin cue to signify its anti-inflammatory and anti-oxidant capacity (GSH, MDA). Nevertheless, co-administration of OLM to PIO showed a synergistic improvement in all the aforementioned parameters in a dose dependent manner. Additionally, OLM with PIO10 provoked a surge in hepatic PPAR γ and adiponectin (5 and 6 folds) with a sharp decrease of about 85% in the NF- κ B/IL-6/p-STAT3/SCOS3 pathway. These effects were confirmed by the histopathological study. In conclusion, OLM and its combination with PIO enhanced insulin sensitivity and guarded against hepatic complications associated with type 2 diabetes probably via modulating various inter-related pathways; namely, metabolic alteration, renin-angiotensin system, inflammatory trajectories, as well as oxidative stress. This study manifests the potential synergistic effects of OLM as an adjuvant therapy to the conventional antidiabetic therapies.

European Journal of Pharmacology 2020, May