Polyphenolic Profile And Biological Activity Of Salvia Splendens Leaves

Wafaa Moustafa El Kady ,Fatma Abd-elkader Moharram, Mohamed Soubhi Marzouk, Siham Mustafa El-Shenawy and Wafaa Mostafa El Kady

Abstract

Objectives: The aim of this study was to investigate a new flavone triglycoside, together with eleven phenolic metabolites from 80% aqueous methanol extract of S. Splendens leaves (AME) and assessment of its hypoglycemic and anti-inflammatory activities along with in vitro antioxidant effect.

Methods: The phenolic composition of S. Splendens leaves was analyzed using UV, 1D and 2D NMR and negative ESI-MS spectroscopy. Hypoglycemic activity of AME was assessed by measuring blood glucose in streptozotocin induced diabetic rats. Key findings: Twelve phenolic metabolites including three phenolic acids, namely cafeeic acid 1, rosmarinic acid 2 and methyl rosmarinate 3, four flavone glycosides viz the new compound luteolin. Moreover, AME exhibited a significant anti-inflammatory activity only at 100 mg/kg in comparison to indomethacin. Finally, AME exhibited a marked significant scavenging activity against DPPPH, the maximum reactive reaction rate after 5 min was 62.9, 82.5, 83.7, 84.3 and 85.1 % for the concentration 10, 20, 30, 40 and 50 mg/ml, respectively. Conclusions: This is the first study reporting the identification of a new flavone along with eleven known phenolic metabolites from AME of S. Splendens. It showed significant hypoglycemic and anti-inflammatory effects in dose dependent manner. Moreover, it showed an in vitro antioxidant activity.

The Journal of pharmacy and pharmacology 2012, January