

Production, characterization and bioinformatics analysis of l-asparaginase from a new *Stenotrophomonas maltophilia* EMCC2297 soil isolate

Nada Anwar AbdelRazek , Walid F. Elkhatib , Marwa M. Raafat and Mohammad M. Aboulwafa

Abstract

An exhaustive screening program was applied for scoring a promising l-asparaginase producing-isolate. The recovered isolate was identified biochemically and molecularly and its l-asparaginase productivity was optimized experimentally and by Response Surface Methodology. The produced enzyme was characterized experimentally for its catalytic properties and by bioinformatics analysis for its immunogenicity. The promising l-asparaginase producing-isolate was selected from 722 recovered isolates and identified as *Stenotrophomonas maltophilia* and deposited at Microbiological Resources Centre (Cairo Mircen) under the code EMCC2297. This isolate produces both intracellular (type I) and extracellular (type II) l-asparaginases with about 4.7 fold higher extracellular l-asparaginase productivity. Bioinformatics analysis revealed clustering of *Stenotrophomonas maltophilia* l-asparaginase with those of *Pseudomonas* species and considerable closeness to the two commercially available l-asparaginases of *E. coli* and *Erwinia chrysanthemi*.

Fourteen antigenic regions are predicted for *Stenotrophomonas maltophilia* l-asparaginase versus 16 and 18 antigenic regions for the *Erwinia chrysanthemi* and *E. coli* l-asparaginases. Type II l-asparaginase productivity of the test isolate reached 4.7 IU/ml/h and exhibited maximum activity with no metal ion requirement at 37°C, 8.6, 40 mM asparagine concentration and could tolerate NaCl concentration up to 500 mM and retain residual activity of 55% at 70°C after half an hour treatment period. Application both of random mutation by gamma irradiation and Response Surface Methodology that determined 38.11°C, 6.89 pH, 19.85 h and 179.15 rpm as optimum process parameters could improve the isolate l-asparaginase productivity. Maximum production of about 8 IU/ml/h was obtained with 0.4% dextrose, 0.1% yeast extract and 10 mM magnesium sulphate. In conclusion l-asparaginase of the recovered *Stenotrophomonas maltophilia* EMCC2297 isolate has characters enabling it to be used for medical therapeutic application

AMB Express 2020, April